arXiv:1304.1614v1 [math.GT] 5 Apr 2013

Some examples of dynamically defined ambient homogeneous wild knots in higher dimensions *

Gabriela Hinojosa[†], Alberto Verjovsky[‡]

April 4, 2013

Dedicated to Francisco González Acuña,
Fico, on the occasion of his 70^{th} birthday.

Abstract

In this paper we consider the Kleinian groups acting conformally on the sphere $\mathbb{S}^{n+2}$ $(1 \leq n \leq 5)$ which have as limit sets wild spheres K^n which were constructed in [4] and prove that K^n is ambient homogeneous. In other words, given two points $p,\ q \in K$ there exists a homeomorphism $\psi : \mathbb{S}^{n+2} \to \mathbb{S}^{n+2}$ such that $\psi(K) = K$ and $\psi(p) = q$.

1 Introduction

In 1920's Antoine [1] and Alexander [2] exhibited embeddings of $\mathbb{S}^2$ in $\mathbb{S}^3$ which are not flat (wild embeddings). These examples can be modified to produce wild embeddings of $\mathbb{S}^n$ in $\mathbb{S}^{n+1}$, for $n \geq 2$.

In general, wild knots behave very different from tame knots. For instance, the wild knot A given by Artin and Fox (see Figure 1), has a wild point p. This knot is not only an example of a wild knot but it is also an example of a knot which is not homogeneous.

We recall that a knot $K \subset \mathbb{S}^{n+2}$ is *ambient homogeneous* if given two points $p,\ q \in K$, there exists a homeomorphism $\psi : \mathbb{S}^{n+2} \to \mathbb{S}^{n+2}$ such that

---

[*] *AMS Subject Classification.* Primary: 57M30. Secondary: 57M45, 57Q45, 30F40. *Key Words.* Wild knots and Kleinian Groups.

[†] This work was partially supported by CONACyT (Mexico), CB-2009-129939

[‡] This work was partially supported by CONACyT (Mexico), CB-2009-129280 and PAPIIT (Universidad Nacional Autónoma de México) #IN100811.



Figure 1: *An example of a wild knot with one wild point.*

$\psi(K) = K$ and $\psi(p) = q$.

Since A contains just one wild point p, it is not possible to give a homeomorphism $\psi : \mathbb{S}^3 \to \mathbb{S}^3$ such that $\psi(K) = K$ and $\psi(p) = q$, $q \neq p$, because any homeomorphism sends wild points into wild points. Notice that any tame knot is homogeneous.

In this paper, we will prove that wild knots of dimensions 1 to 5, which are obtained as limit sets of some Kleinian groups (dynamically defined wild knots) are homogeneous (see Corollary 4.2).

2 Preliminaries

We will give some basic definitions of Kleinian groups.

Let $M\ddot{o}b(\mathbb{S}^n)$ denote the group of Möbius transformations of the n-sphere $\mathbb{S}^n = \mathbb{R}^n \cup \{\infty\}$ *i.e.*, the group of diffeomorphisms of the n-sphere that consist of composition of inversions in $\mathbb{S}^n$. The group $M\ddot{o}b(\mathbb{S}^n)$ admits an extension to the hyperbolic space $\mathbb{H}^{n+1}$, so that we can identify the group $M\ddot{o}b(\mathbb{S}^n)$ with the group of isometries of hyperbolic $(n+1)$-space $\mathbb{H}^{n+1}$.

Definition 2.1 *A discrete subgroup* $\Gamma \subset M\ddot{o}b(\mathbb{S}^n)$ *is called a Kleinian group.*

The *discontinuity set* $\Omega(\Gamma)$ of a group $\Gamma \subset M\ddot{o}b(\mathbb{S}^n)$, is the largest open subset in $\mathbb{S}^n$ where Γ acts properly discontinuously. Its complement $\mathbb{S}^n \setminus \Omega(\Gamma)$ is the limit set $\Lambda(\Gamma)$ of the group Γ. Equivalently, the limit set of a Kleinian group can be described as the accumulation set in the sphere $\mathbb{S}^n$ of an orbit Γx; where x is an arbitrary point in $\mathbb{H}^{n+1}$.

One way to illustrate the action of a Kleinian group Γ is to draw a picture of $\Omega(\Gamma)/\Gamma$. For this purpose a fundamental domain is very helpful.

Definition 2.2 *A fundamental domain D for a Kleinian group Γ is a codimension zero piecewise-smooth submanifold (sub polyhedron) of $\Omega(\Gamma)$ satisfying the following*

1. $\bigcup_{g\in\Gamma} g(Cl_{\Omega(\Gamma)}D) = \Omega$ *where* $Cl_{\Omega(\Gamma)}$ *is the closure in* $\Omega(\Gamma)$.
2. $g(Int(D)) \cap Int(D) = \emptyset$ *for all* $g \in \Gamma \setminus \{e\}$, *where e is the identity in Γ and Int denotes the interior.*
3. *The boundary of D in $\Omega(\Gamma)$ is a piecewise-smooth (polyhedron) submanifold in $\Omega(\Gamma)$, divided into a union of smooth submanifolds (convex polyhedra) which are called* faces. *For each face S, there is a corresponding face F and an element $g_{SF} \in \Gamma \setminus \{e\}$ such that $g_{SF}(S) = F$ (g_{SF} is called a face-pairing transformation); $g_{SF} = g_{FS}^{-1}$.*
4. *Only finitely many translates of D meet any compact subset of $\Omega(\Gamma)$.*

THEOREM 2.3 *([8], [9]) Let $D^* = \overline{D} \cap \Omega / \sim_G$ denote the orbit space with the quotient topology. Then D^* is homeomorphic to Ω/G.*

3 Criterion of homogeneity

Next, we will establish a criterion of homogeneity. This criterion is elemental, however we would like to emphasize that our main difficulty lies on proving that dynamically defined wild knots (see next section) satisfy it.

THEOREM 3.1 (Homogeneity criterion). *Let $\varphi_n : \mathbb{D}^2 \times \mathbb{S}^n \to \mathbb{S}^{n+2} = \mathbb{R}^{n+2} \cup \{\infty\}$, $n = 1, 2, \ldots$ be differentiable embeddings satisfying the following.*

1. $\mathrm{Im}(\varphi_n) = T_n$
2. $T_{n+1} \subset \mathrm{Int}(T_n)$
3. $\mathrm{diam}(\varphi_n(\mathbb{D}^2 \times \{x\})) \to 0$ *as* $n \to \infty$ *independently of* $x \in \mathbb{S}^n$.
4. $\Sigma = \cap T_n \cong \mathbb{S}^n$ *(possibly fractal).*
5. *Let $z \in \partial\mathbb{D}^2$ be a fixed point. The sequence of longitude manifolds $h_n(x) = \varphi_n(z, x) \subset T_n$ converges to a parametrization of Σ, i.e. an embedding $h : \mathbb{S}^n \to \mathbb{R}^{n+2}$ such that $h(\mathbb{S}^n) = \Sigma$.*

Then Σ is ambient homogeneous.

Proof. Let $x,\ y \in \Sigma$. So $x = h(x_1)$ and $y = h(y_1)$, where $x_1,\ y_1 \in \mathbb{S}^n$. Let $A \in SO(n+2)$ such that $Ax_1 = y_1$ and define the rotation map $g : \mathbb{S}^n \to \mathbb{S}^n$ as $g(z) = Az$, $z \in \mathbb{S}^n$. Consider the map $f : \Sigma \to \Sigma$ given by $f = h \circ g \circ h^{-1}$, thus f is a homeomorphism such that $f(x) = y$.

By the above, we have a sequence of homeomorphisms $\alpha_n : \partial T_n \to \partial T_n$ given by $\alpha_n = \varphi_n \circ \eta \circ \varphi_n^{-1}$, where $\eta = \mathrm{Id} \times g : \mathbb{D}^2 \times \mathbb{S}^n \to \mathbb{D}^2 \times \mathbb{S}^n$. In particular, we have a sequence of longitude manifold homeomorphisms $f_n = h_n \circ \eta \circ h_n^{-1}$ such that $\lim_{n\to\infty} f_n = \lim_{n\to\infty} h_n \circ \eta \circ h_n^{-1} = f$.

We define the homeomorphism $\psi_0 : (\mathbb{S}^{n+2} - \mathrm{Int}(T_1)) \to (\mathbb{S}^{n+2} - \mathrm{Int}(T_1))$ in such a way that $\psi_0|_{\partial T_1} = \alpha_1$. Now, we consider the homeomorphism $\mathcal{K}_1 : (\partial T_1 \cup \partial T_2) \to (\partial T_1 \cup \partial T_2)$ given by $\mathcal{K}_1|_{\partial T_1} = \psi_0$ and $\mathcal{K}_2|_{\partial T_2} = \alpha_2$. Observe that each $\mathcal{K}_1$ is isotopic to the identity map, then we can apply the next lemma to extend it to a homeomorphism $\psi_1 : (T_1 - \mathrm{Int}(T_2)) \to (T_1 - \mathrm{Int}(T_2))$.

LEMMA 3.2 *Let M be a compact $(n+2)$-manifold with $\partial M \neq \emptyset$ (no necessarily connected). Let $g : \partial M \to \partial M$ be a homeomorphism which is isotopic to the identity map $I_{\partial M}$. Then g admits an extension $G : M \to M$.*

Proof. Let $\psi : \partial M \times [0,1] \to M$ be a collaring of the boundary, i.e., ψ is a homeomorphism such that $\psi(x,0) = x$. Let $N = \psi(\partial M \times [0,1]) \subset M$. Let $\{g_t\},\ t \in [0,1]$, be an isotopy of g to the identity, i.e., $g_o = g$, $g_1 = I_{\partial M}$. Let $H : \partial M \times [0,1] \to \partial M \times [0,1]$ be given by the formula $H(x,t) = (g_t(x), t)$. Let $G_0 : N \to N = \psi \circ H \circ \psi^{-1}$. Then we define $G : M \to M$ as $G(y) = G_0(y)$ if $y \in N$ and $G(y) = y$ if $y \notin N$. □

We continue inductively, so at the m-stage we have a homeomorphism $\psi_m : (T_m - \mathrm{Int}(T_{m+1})) \to (T_m - \mathrm{Int}(T_{m+1}))$, such that $\psi_m|_{\partial T_{m+1}} = \alpha_{m+1}$ and $\psi_m|_{\partial T_m} = \psi_{m-1}$.

This construction allows us to define a map $\psi : (\mathbb{S}^{n+2} - \Sigma) \to (\mathbb{S}^{n+2} - \Sigma)$ as $\psi(x) = \psi_m(x)$ if $x \in (T_m - \mathrm{Int}(T_{m+1}))$. Notice that ψ is a homeomorphism, since each ψ_m is a homeomorphism and $\psi_m(x) = \psi_{m+1}(x)$ for $x \in \partial T_{m+1}$ and for all m. Now, using that $\Sigma = \cap T_n \cong \mathbb{S}^n$ and $\lim_{n\to\infty} f_n = \lim_{n\to\infty} h_n \circ \eta \circ h_n^{-1} = f$, we extend ψ by continuity to a homeomorphism $\tilde{\psi} : \mathbb{S}^{n+2} \to \mathbb{S}^{n+2}$, such that $\tilde{\psi}|_\Sigma = f$; hence $\tilde{\psi}(x) = y$. Therefore Σ is ambient homogeneous. □

4 Dynamically defined wild knots

In this section, we will prove that dynamically defined wild knots are ambient homogeneous. We will start describing them.

4.1 Dynamically defined wild knots in higher dimensions

We will give a brief description of the construction of dynamically defined wild knots in higher dimensions (for more details see [4]).

Let K_1 be a smooth fibered knot of dimension n in $\mathbb{S}^{n+2}$. In [4] was proved that for $n = 1, 2, 3, 4$ and 5, there exists a finite collection of $(n+2)$-dimensional closed round balls, B_1, B_2, $\ldots$, B_r such that the balls B_i and B_j are either disjoint, tangent or their boundaries meet orthogonally, and satisfies that $K_1 \subset \cup_{i=1}^{r} B_i$ and the limit set of the group Γ generated by reflections on its boundaries is a *wild* n-sphere embedded in the $(n+2)$-sphere.

Let $|T_1| = \cup_{j=1}^{r} B_j$. Notice that it has some points of tangency c_i, $i = 1, \ldots, d$, then we set round closed balls B_{C_i} centered at c_i. The union $|\tilde{T_1}| = \cup_{i=1}^{d} B_{C_i} \cup |T_1|$ is called a *generalized necklace*, and it is a closed regular neighborhood of K_1 (corollary 4.6 in [4]).

In general, in order to obtain a wild knot in high dimensions as the limit set of a geometrically finite Kleinian group we need to construct a finite collection of balls satisfying the above properties.

We will describe the limit set of Γ. Recall that Γ is generated by reflections I_j on ∂B_j, $j = 1, 2, \ldots, r$ and to find its limit set, we need to find all the accumulation points of its orbits. We will do this by stages.

Stage I. We will apply induction on the number of reflections.

1. *First step*: We reflect with respect to ∂B_1, *i.e.*, we apply $I_1 \in \Gamma$. The set $D_1^{'} = (|\tilde{T_1}| - B_1) \cup I_1(|\tilde{T_1}|)$ is a regular neighborhood of the geometric realization of its nerve, which is isotopic to the connected sum of K_1 with its mirror image $-K_1$ (see [4]).

2. *Second step*: We reflect with respect to ∂B_2, *i.e.*, we apply $I_2 \in \Gamma$. The set $D_2^{'} = (|\tilde{T_1}| - \cup_{i=1}^{2} B_i) \cup I_1(|\tilde{T_1}|) \cup I_2(D_1^{'})$ is a regular neighborhood of the geometric realization of its nerve, which is isotopic to the connected sum $K_1 \# (-K_1) \# K_1 \# (-K_1)$.

3. r^{th}-*step*: We reflect with respect to ∂B_r, *i.e.*, we apply $I_r \in \Gamma$. The set $D_r^{'} = I_1(|\tilde{T}_1|) \cup I_2(D_1^{'}) \cup I_3(D_2^{'}) \cup \ldots \cup I_r(D_{r-1}^{'})$ is a regular neighborhood of the geometric realization of its nerve, which is isotopic to the connected sum of 2^{r-1} copies of K_1 and 2^{r-1} copies of $-K_1$.

At the end of the r^{th}-step , we obtain a regular neighborhood $|\tilde{T}_2|$ of a new tame n-knot K_2, which is isotopic to the connected sum of 2^{r-1} copies of K_1 and 2^{r-1} copies of $-K_1$. Notice that $|\tilde{T}_2| \subset \mathrm{Int}(|\tilde{T}_1|)$.

Stage II.

Repeat k-times Stage I. At this stage we obtain a regular neighborhood $|\tilde{T}_{k+1}|$ of a new tame n-knot K_{k+1} which is isotopic to the connected sum of 2^{kr-1} copies of K_1 and 2^{kr-1} copies of $-K_1$. By construction, $|\tilde{T}_{k+1}| \subset \mathrm{Int}(|\tilde{T}_k|)$.

The limit set of Γ is given by

$$\Lambda(\Gamma) = \varprojlim_{l} |\tilde{T}_l| = \bigcap_{l=1}^{\infty} |\tilde{T}_l|.$$

It has been proved (see [4]) that the limit set $\Lambda(\Gamma)$ is an n-knot. If we start with a *nontrivial* fibered knot K_1, then its limit set is a wild knot whose complement fibers over the circle where the fibre has infinitely generated homology groups. The knot is called a *dynamically-defined fibered wild knot.*

4.2 Homogeneity

We will prove the following result using the homogeneity criterion 3.1.

THEOREM 4.1 *Dynamically-defined wild knots are ambient homogeneous.*

Proof. Let K_1 be a smooth n-knot and $|\tilde{T}_1|$ be a generalized necklace subordinated to K_1. Observe that by construction we have a sequence of closed tubular neighborhoods $\{|\tilde{T}_k|\}_{k=1}^{\infty}$ such that $|\tilde{T}_k| \subset \mathrm{Int}(|\tilde{T}_{k-1}|)$ and $\Lambda(\Gamma) = \varprojlim_{l} |\tilde{T}_l| = \bigcap_{l=1}^{\infty} |\tilde{T}_l|$. So, we have that dynamically-defined wild knots satisfy the conditions 1 to 4 of theorem 3.1.

Suppose that $|\tilde{T}_1| = \cup_{j=1}^{m_1} B_j$, where B_j is a round $(n+2)$-closed ball. Let $\pi_1 : |\tilde{T}_1| \to K$ be the projection. Now, we will subdivide $\mathbb{S}^n$ into connected closed sets A_i, for each $i = 1, \ldots, m_1$; following the same pattern that $|\tilde{T}_1| \cap K_1$, *i.e.* $A_i \neq \emptyset$ corresponds to the closed set $V_i = B_i \cap K_1$ and $A_j \cap A_l \neq \emptyset$ if and only if $V_j \cap V_l \neq \emptyset$. Moreover, let $\phi_1 : \mathbb{D}^2_{r_1} \times \mathbb{S}^n \to |\tilde{T}_1|$ be a homeomorphism. We can assume, up to reparametrization, that $\phi_1(\mathbb{D}^2_{r_1}, A_i) = \pi_1^{-1}(V_i)$ and $\phi_1(\mathbb{D}^2_{r_1}, A_i \cap A_j) = \pi_1^{-1}(V_i \cap V_j)$.

We define the longitude map $h_1 : \mathbb{S}^n \to |\tilde{T}_1|$ as $h_1(x) = \phi_1(r_1 z \times x)$, for a fixed point $z \in \partial \mathbb{D}^2$ and $x \in \mathbb{S}^n$.

For the second stage on the reflecting process, we have a generalized necklace $|\tilde{T}_2| = \cup_{j=1}^{m_2} B_j^2$ subordinate to the knot K_2. Let $\pi_2 : |\tilde{T}_2| \to K_2$ be the projection. Now, we will subdivide each closed set A_i of $\mathbb{S}^n$ into connected closed sets A_{ij} following the same pattern that $B_i \cap |\tilde{T}_2| \cap K_2$, *i.e.* $A_{ij} \neq \emptyset$ corresponds to the closed set $V_{ij} = B_i \cap B_j^2 \cap K_2 \neq \emptyset$ and $A_{ij} \cap A_{ls} \neq \emptyset$ if and only if $V_{ij} \cap V_{ls} \neq \emptyset$. Let $\phi_2 : \mathbb{D}^2_{r_2} \times \mathbb{S}^n \to |\tilde{T}_2|$ be a homeomorphism. We can assume, up to reparametrization, that $\phi_2(\mathbb{D}^2_{r_2}, A_{ij}) = \pi_2^{-1}(V_{ij})$ and $\phi_2(\mathbb{D}^2_{r_2}, A_{ij} \cap A_{ls}) = \pi_2^{-1}(V_{ij} \cap V_{ls})$. Notice that $\pi_2^{-1}(V_{ij}) \subset \pi_1^{-1}(V_i)$. Now, we consider the corresponding longitude map $h_2(x) = \phi_2(r_2 z \times x)$ given by $h_2(x) = \phi_1(r_2 z \times x)$, $x \in \mathbb{S}^n$.

We continue inductively. So at the m-stage of the reflecting process, we have a generalized necklace $|\tilde{T}_m| = \cup_{j=1}^{m_m} B_j^m$ subordinate to the knot K_m. Let $\pi_m : |\tilde{T}_m| \to K_m$ be the projection. Now, we subdivide each connected closed set $A_{1_{k_1},2_{k_2},\ldots,(m-1)_{k_{(m-1)}}}$ into connected closed sets $A_{1_{k_1},2_{k_2},\ldots,m_{k_m}}$ following the same pattern that $\pi_{k-1}^{-1}(V_{1_{k_1},2_{k_2},\ldots,(m-1)_{k_{(m-1)}}}) \cap |\tilde{T}_m|$, in other words, the closed set $A_{1_{k_1},2_{k_2},\ldots,m_{k_m}} \neq \emptyset$ corresponds to the closed set $V_{1_{k_1},2_{k_2},\ldots,m_{k_m}} = \pi_{k-1}^{-1}(V_{1_{k_1},2_{k_2},\ldots,(m-1)_{k_{(m-1)}}}) \cap B_{k_m}^{m_m} \cap K_m \neq \emptyset$.

Let $\phi_m : \mathbb{D}^2_{r_m} \times \mathbb{S}^n \to |\tilde{T}_m|$ be a homeomorphism. We can assume, up to reparametrization, that $\phi_m(\mathbb{D}^2_{r_m}, A_{1_{k_1},2_{k_2},\ldots,m_{k_m}}) = \pi_m^{-1}(V_{1_{k_1},2_{k_2},\ldots,m_{k_m}})$ and $\phi_m(\mathbb{D}^2_{r_m}, A_{1_{k_1},2_{k_2},\ldots,m_{k_m}} \cap A_{1_{l_1},2_{l_2},\ldots,m_{l_m}}) = \pi_m^{-1}(V_{1_{k_1},2_{k_2},\ldots,m_{k_m}} \cap V_{1_{l_1},2_{l_2},\ldots,m_{l_m}})$. By construction, $\pi_m^{-1}(V_{1_{k_1},2_{k_2},\ldots,m_{k_m}}) \subset \pi_{k-1}^{-1}(V_{1_{k_1},2_{k_2},\ldots,(m-1)_{k_{(m-1)}}})$. Consider the longitude map $h_m(x) = \phi_m(r_m z \times x)$, $x \in \mathbb{S}^n$.

Let $h(x) = \lim_{m \to \infty} h_m(x)$, $x \in \mathbb{S}^n$. Then $h : \mathbb{S}^n \to \mathbb{S}^{n+2}$ is a continuous map. In fact, for each $x \in \mathbb{S}^n$ there exists, by construction, a nested sequence of closed sets $\{A_{1_{k_1},2_{k_2},\ldots,m_{k_m}}\}$ such that $x = \cap_{m=1}^{\infty} A_{1_{k_1},2_{k_2},\ldots,m_{k_m}}$. Notice that this sequence converges to x with respect to the Hausdorff metric of closed sets on $\mathbb{S}^n$. Now $h_m(x) \in \pi_m^{-1}(V_{1_{k_1},2_{k_2},\ldots,m_{k_m}})$, so $h(x) \in$

$\pi_m^{-1}(V_{1_{k_1},2_{k_2},\ldots,m_{k_m}}) \subset |\tilde{T}_m|$; but $\{\pi_m^{-1}(V_{1_{k_1},2_{k_2},\ldots,m_{k_m}})\}$ is a nested sequence of closed sets; hence it converges to $h(x) = \cap_{m=1}^{\infty}\pi_m^{-1}(V_{1_{k_1},2_{k_2},\ldots,m_{k_m}}) \in \Lambda$ with respect to the Hausdorff metric of closed sets on $\mathbb{S}^{n+2}$.

Observe that for each $p \in \Lambda$, there exists a nested sequence of closed sets $\{\pi_m^{-1}(V_{1_{j_1},2_{j_2},\ldots,m_{j_m}})\}$, where $\pi_m^{-1}(V_{1_{j_1},2_{j_2},\ldots,m_{j_m}}) \subset |\tilde{T}_m|$ and such that $p = \cap_{m=1}^{\infty}\pi_m^{-1}(V_{1_{j_1},2_{j_2},\ldots,m_{j_m}})$. This implies that h is onto. Moreover, given $h(x_1) = h(x_2) = p \in \Lambda$, since $p = \cap_{m=1}^{\infty}\pi_m^{-1}(V_{1_{j_1},2_{j_2},\ldots,m_{j_m}})$ we have that $h_m(x_1), h_m(x_2) \in \pi_m^{-1}(V_{1_{j_1},2_{j_2},\ldots,m_{j_m}})$, for all m. Thus x_1, $x_2 \in A_{1_{k_1},2_{k_2},\ldots,m_{k_m}}$ for all m. This implies that x_1, $x_2 \in \cap_{m=1}^{\infty}A_{1_{k_1},2_{k_2},\ldots,m_{k_m}}$. Therefore, $x_1 = x_2$. As a consequence, $h : \mathbb{S}^n \to \mathbb{S}^{n+2}$ is a parametrization of Λ. Applying theorem 3.1, we have that Λ is ambient homogeneous. $\square$

COROLLARY 4.2 *The infinite family of dynamically defined wild knots* $K^n \hookrightarrow \mathbb{S}^{n+2}$, $1 \leq n \leq 5$, *constructed in [4], are ambient homogeneous.*

References

[1] L. Antoine, *Sur l'homeomorphiede deux figures et de leurs voisinages*, J. Math. Pures. Appl., 86 (1921), 221-235.

[2] J. W. Alexander, *On the subdivision of 3-space by a polyhedron*, Proc. Nat. Acad. Sci. U.S.A., 10 (1924), 6-8.

[3] E. Artin, *Zur Isotopie zweidimensionalen Flächen im* R_4 Abh. Math. Sem. Univ. Hamburg (1926), 174–177.

[4] M. Boege, G. Hinojosa, and A. Verjovsky. *Wild knots in higher dimensions as limit sets of Kleinian groups.* Conform. Geom. Dyn. 13 (2009), 197–216.

[5] R. H. Fox. *A Quick Trip Through Knot Theory.* Topology of 3-Manifolds and Related Topics. Prentice-Hall, Inc., 1962..

[6] G. Hinojosa. *Wild knots as limit sets of Kleinian Groups.* Contemporary Mathematics 389 (2005), 125–139.

[7] G. Hinojosa, A. Verjovsky *Homogeneity of dynamically defined wild knots.* Rev. Mat. Compl. vol. 19 no. 1, 2006. Pags. 101-111.

[8] M. Kapovich. *Hyperbolic Manifolds and Discrete Groups.* Progress in Mathematics, Birkhauser, 2001.

[9] B. Maskit. *Kleinian Groups.* Springer Verlag, 1987.

[10] D. Rolfsen. *Knots and Links.* Publish or Perish, Inc. 1976.

[11] B. Rushing. *Topological Embeddings.* Academic Press, 1973, Vol 52.

G. Hinojosa. Universidad Autónoma del Estado de Morelos. Av. Universidad 1001, Col. Chamilpa. Cuernavaca, Morelos, México, 62209.
E-mail address: gabriela@uaem.mx

A. Verjovsky. Instituto de Matemáticas, Universidad Nacional Autónoma de México, Unidad Cuernavaca. Av. Universidad s/n, Col. Lomas de Chamilpa. Cuernavaca, Morelos México, 62209.
E-mail address: alberto@matcuer.unam.mx